

Importance of dietary iron for women

References

1. Australian Bureau of Statistics, 2015 *Australian Health Survey: Usual Nutrient Intakes* [Australian Health Survey: Usual Nutrient Intakes, 2011-12 financial year | Australian Bureau of Statistics (abs.gov.au)] Accessed 9 March 2022.
2. National Health and Medical Research Council and New Zealand Ministry of Health, 2006, *Nutrient Reference Values for Australia and New Zealand* <https://www.nrv.gov.au/nutrients/iron>, accessed 9 March 2022.
3. Ahmed F, Coyne T, Dobson A, McClintock C. Iron status among Australian adults: findings of a population-based study in Queensland, Australia. *Asia Pac J Clin Nutr* 2008;17:40-7.
4. Greig A et al. Iron deficiency, cognition, mental health and fatigue in women of childbearing age: a systematic review *Journal of Nutritional Science* 2013;2:E14. doi:10.1017/jns.2013.7.
5. Lomagno KA et al. Increasing iron and zinc in pre-menopausal women and its effects on mood and cognition: a systematic review. *Nutrients* 2014; 6:5117-41. doi:10.3390/nu115117.
6. Pasricha SR, Flecknoe-Brown SC, Allen KJ, Gibson PR, McMahon LP, Olynyk JK, et al. Diagnosis and management of iron deficiency anaemia: a clinical update. *Med J Aust* 2010;193:525-32.
7. Australian Bureau of Statistics, 2014, *Australian Health Survey: Nutrition First Results - Foods and Nutrients, 2011-12*, 'Table 10: Proportion of Nutrients from food groups', data cube: Excel spreadsheet, cat. no. 4364.0.55.007.
8. Heath A, Skeaff C, O'Brien S, Williams S, Gibson R. Can dietary treatment of non-anemic iron deficiency improve iron status? *J Am Coll Nutr* 2001;20(5): 477-484. doi:10.1080/07315724.2001.10719056.
9. Patterson A, Brown W, Roberts D, Seldon M. Dietary treatment of iron deficiency in women of childbearing age. *Am J Clin Nutr* 2001;74(5):650-56. doi:10.1093/ajcn/74.5.650.
10. Cheng H, Griffin H, Byrant C, Rooney K, Steinbeck K, O'Connor H. Impact of diet and weight loss on iron and zinc status in overweight and obese young women. *Asia Pac J Nutr* 2013;22(4):574-82. doi: 10.6133/apjcn.2013.22.4.08.
11. O'Connor H, Munas Z, Giffin H, Rooney K, Cheng HL, Steinbeck K Nutritional adequacy of energy restricted diets for young obese women. *Asia Pacific Journal of Clinical Nutrition* 2011;20:206-211.
12. National Health and Medical Research Council. *Australian Dietary Guidelines* [Internet]. Canberra (ACT): NHMRC; 2013 [cited 2022 March 9]. Available from: <https://www.nhmrc.gov.au/adg>
13. Sui Z, Raubenheimer D, Rangan A. Consumption patterns of meat, poultry, and fish after disaggregation of mixed dishes: secondary analysis of the Australian National Nutrition and Physical Activity Survey 2011–12. *BMC Nutr* 2017;3:52. doi:10.1186/s40795-017-0171-1.
14. Fayet F, Flood V, Petocz P, Samman S. Avoidance of meat and poultry decreases intakes of omega-3 fatty acids, vitamin B12, selenium and zinc in young women. *J Hum Nutr Diet* 2014;27:135-142. doi:10.1111/jhn.12092.
15. Leonard AJ et al. The effect of nutrition knowledge and dietary iron intake on iron status in young women. *Appetite*; 2014;81:225–231.
16. Lim K, Booth A, Szymlek-Gay EA, Gibson RS, Bailey KB, Irving D, Nowson C, Riddell L Associations between dietary iron and zinc intakes, and between biochemical iron and zinc status in women. *Nutrients* 2015;7:2983-99. doi.org/10.3390/nu7042983.

17. Food and Agriculture Organization of the United Nations & World Health Organization
Requirements of vitamin A, iron, folate and vitamin B12: report of a joint FAO/WHO expert consultation. Rome:Lanham MD: FAO; UNIPUB [distributor]; 1988.
18. Schönfeldt H and Hall N. Determining iron bioavailability with a constant heme iron value.
Journal of Food Composition and Analysis 2011;24:738-740.